arXiv:1210.1156v1 [math.PR] 3 Oct 2012

Local Malliavin Calculus for Lévy Processes and Applications [*]

Jorge A. León[†]
Departamento de Control Automático
Cinvestav-IPN
Apartado Postal 14-740
07000 México D.F., Mexico
jleon@ctrl.cinvestav.mx

Josep L. Solé, Frederic Utzet
Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193-Cerdanyola (Catalunya), Spain
jllsole@mat.uab.cat
utzet@mat.uab.cat

Josep Vives
Departament de Probabilitat, Lògica i Estadística
Universitat de Barcelona
08007-Barcelona (Catalunya), Spain
josep.vives@ub.edu

08-03-2012

Abstract

In this paper a Malliavin calculus for Lévy processes based on a family of true derivative operators is developed. The starting point is an extension to Lévy processes of the pioneering paper by Carlen and Pardoux [8] for the Poisson process, and our approach includes also the classical Malliavin derivative for Gaussian processes. We obtain a sufficient condition for the absolute continuity of functionals of the Lévy process. As an application, we analyze the absolute continuity of the law of the solution of some stochastic differential equations.

Keywords: Lévy process; Malliavin derivative; stochastic differential equations
AMS 1991 subject classifications: 60H05, 60H10

1 Introduction

Over the past twenty five years, it has been done several attempts for extending the Malliavin calculus to a scenario driven by processes with jumps. Among these efforts, we should cite the celebrated papers of Bismut [6], Léandre [18] and the monograph of Bichteler, Gravereaux and Jacod [5]. In the latter it is assumed the restriction that the Lévy measure ν is absolutely continuous with respect to the Lebesgue measure and the main idea is to derive with respect to the size of the jumps. However, if

---

[*] Partially supported by the Ministerio de Ciencia e Inovación grant MTM2009-08869 and FEDER
[†] Partially supported by the CONACyT grant 98998

the measure ν is discrete, the condition in [5] is not satisfied, and by this reason Carlen and Pardoux [8] proposed, for a Poisson process, a derivative operator based on translation of the jump times.

There is another approach to define a Malliavin derivative as an annihilation operator through a chaos expansion. In this way, Nualart and Vives [23] proved that for the standard Poisson process the annihilation operator (Malliavin derivative) agrees with a translation operator. Immediately it appeared the necessity of extending these methods to Lévy processes with a more general Lévy measure, and many authors proposed and studied such difference operator: see for example, Picard [24] and [25], members of the norwegian school of probabilities like Di Nunno, Meyer-Brandis, Øksendal and Proske [13] and Benth and Lokka [4], and also Ishikawa and Kunita [15], and Solé, Utzet and Vives [26]. It is also interesting to cite the extension to abstract Fock spaces by Applebaum [2]. Note that in all those works, in contrast with the Wiener case, the annihilation operator is not a true derivative and it does not satisfy the chain rule.

A number of authors have made contributions in the direction of defining a true derivation operator for jump processes, under more or less restrictive assumptions. We should cite Denis [11] and Bouleau and Denis [7], who use the context of Dirichlet forms, Di Nunno [12], and recently Kulik [17] and Bally [3]. We have to note that Decreusefond and Savy [10] worked with marked point processes (i.e., processes with compensator $\lambda(s)ds\nu(dz))$, introducing an operator similar to the one studied in this paper, but with the restriction that ν should be a probability measure.

In this paper we define a true derivative operator for functionals of a general Lévy processes which can have a continuous part. Such operator is an extension to Lévy processes of the one introduced by Carlen and Pardoux [8] for the Poisson process. Our main finding is that this operator can be restricted to a Borel subset of $\mathbb{R}$, say Λ, where live part of the jump sizes of the Lévy processes, and a suitable (time and space) weight function $k(t, x)$ can be introduced. That subset and weight function can be chosen in every application to suit the properties of the problem in hand. By this reason we call it the *local Malliavin derivative.* Our procedure is, as in the Brownian Malliavin calculus, to start with a class of smooth random variables, define the derivative, and extend by density to a domain contained in $L^2(\Omega)$. We observe that this approach can be used even for the difference operator as it is done in Geiss and Laukkarinen [14], but, as we commented, unfortunately, the chain rule is not satisfied for the difference operator.

We would like to point out that we don't present here a complete Malliavin calculus theory, because our integration by parts formula (Proposition 3.7) does not cover the general case. On the other hand, we do not give any result about the smoothness of the density. However we believe that we may obtain some interesting results from our development.

Using the chain rule of our derivative operator we give a sufficient condition for the absolutely continuity of functionals of the involved Lévy process analogue to the one of Carlen and Pardoux [8]. In particular, we apply that condition to deduce the absolute continuity of solutions of different stochastic differential equations. The first case studied is the equation governed by a Lévy process with continuous part

$$Z_t = x + \int_0^t b(Z_s)ds + \int_0^t \sigma(Z_s)dW_s + \int_0^t \int_{\mathbb{R}_0} l(y)ydN(s, y), \quad t \in [0, T], \tag{1.1}$$

with suitable coefficients b, σ and l, and where $\mathbb{R}_0 := \mathbb{R} - \{0\}$. We get a similar result as the one contained in Nualart [22], where jumps are not considered.

The second application is devoted to the following equation, driven by a pure additive jump process,

$$Z_t = x + \int_0^t f(Z_s)ds + \int_0^t \int_{\mathbb{R}_0} h(y)dN(s, y), \quad t \in [0, T]. \tag{1.2}$$

Here $x \in \mathbb{R}$, $f : \mathbb{R} \to \mathbb{R}$ is a function with a bounded derivative and $h \in L^2(\mathbb{R}_0, \nu) \cap L^1(\mathbb{R}_0, \nu)$, ν is the Lévy measure corresponding to the driving Lévy process. It is well-known that this equation has a unique square-integrable solution. The first step is to find the derivative of the solution at time T. Then, in the case that $\nu(\mathbb{R}_0) < \infty$ and that the function f is monotone, choosing the convenient $k(t, x)$, we obtain that Z_T satisfies the criterion established for the absolute continuity in the set of paths where there is at least one jump. Moreover, if $\nu(\mathbb{R}_0) = \infty$, it is only necessary the condition of monotony for f in a neighborhood of the initial condition x. This is the same type of result obtained by Nourdin and Simon [21] using the stratification method due to Davydov *et al.* [9]. Note that as the Lévy measure is not finite, we have, with probability one, infinity jumps in any neighborhood of the origin, and this fact guarantees that we can find the suitable function k in order to restrict us to the neighborhood of monotony of f.

Finally, we study the equation

$$Z_t = x + \int_0^t f(Z_s)ds + \int_0^t \int_{\mathbb{R}_0} h(y)g(Z_{s-})dN(s, y), \quad t \in [0, T], \tag{1.3}$$

with some conditions on f, g and h. The existence of a density for Z_T was analyzed by Carlen and Pardoux [8] in the case that the involved Lévy process is a Poisson process. When the Lévy process is a compound Poisson, using our criterion of absolute continuity with a suitable function $k(t, x)$ that allows us to concentrate in only one jump, we prove the absolutely continuity of the solution in the set of paths where there is at least one jump in the case where a condition on the Wronskian of f and g is satisfied.

The paper is organized as follows. Section 2 presents the preliminaries and notations. Section 3 is devoted to the study of the family of derivative operators introduced and to the criterion of absolute continuity. In Section 4 we present some properties of the derivative operator applied to multiple integrals which are needed in the next section. Finally, in Section 5 we get the absolute continuity of the solutions to the stochastic differential equations (1.1), (1.2) and (1.3).

2 Preliminaries

In this section we introduce the framework and the notation that we use in this paper. Let $X = \{X_t, t \geq 0\}$ be a Lévy process, that is, X has stationary and independent increments, is continuous in probability, is càdlàg and $X_0 = 0$, with triplet (γ, σ, ν), and it is defined on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$; for all these notions we refer to Sato [27]. By convention, the Lévy measure ν is defined on $\mathbb{R}$ and $\nu(\{0\}) = 0$. Also, let $\{\mathcal{F}_t, t \geq 0\}$ be the natural filtration associated with X. The process X admits the Lévy-Itô representation:

$$X_t = \gamma t + \sigma W_t + \int\int_{(0,t]\times\{|x|>1\}} x dN(s, x) + \lim_{\epsilon \downarrow 0} \int\int_{(0,t]\times\{\epsilon<|x|\leq 1\})} x d\tilde{N}(s, x), \tag{2.1}$$

where W is a standard Brownian motion, N is the jump measure of the process defined below, W and N are independent, and $d\tilde{N}(t, x) = dN(t, x) - dt\nu(dx)$. The limit is a.s., uniform in t in any bounded interval.

For any Borel set $B \in \mathcal{B}((0, \infty) \times \mathbb{R}_0)$, with $\mathbb{R}_0 := \mathbb{R} - \{0\}$, the jump measure is defined by

$$N(B) = \text{card}\{t \in (0, \infty) : (t, \Delta X_t) \in B\},$$

where $\Delta X_t = X_t - X_{t-}$.

It is well known (see Itô [16]) that X determines a centered independent scattered random measure M on $[0,\infty)\times\mathbb{R}$. To this end, consider the measure

$$\mu(dt,dx)=\lambda(dt)\delta_0(dx)+\lambda(dt)x^2\nu(dx),$$

where λ is the Lebesgue measure in $\mathbb{R}$. For $E\in\mathcal{B}([0,\infty)\times\mathbb{R})$ such that $\mu(E)<\infty$,

$$M(E)=\sigma\int_{E(0)}dW_t+\lim_{n\to\infty}\int\int_{\{(t,x)\in E':\frac{1}{n}\leq|x|\leq n\}}xd\tilde{N}(t,x),$$

where $E(0)=\{t>0:(t,0)\in E\}$, $E'=E-\{(t,0)\in E\}$, and the limit is in $L^2(\Omega)$. For $A,B\in\mathcal{B}([0,\infty)\times\mathbb{R}))$, with $\mu(A)<\infty$ and $\mu(B)<\infty$, we have

$$\mathbb{E}[M(A)M(B)]=\mu(A\cap B).$$

Because we work in a finite time interval $[0,T]$, we consider the restriction of M to $[0,T]\times\mathbb{R}$. For a function $h\in L^2([0,T]\times\mathbb{R},\mu)$, we can construct the integral (in the $L^2(\Omega)$ sense)

$$M(h):=\int_{[0,T]\times\mathbb{R}}h(t,x)\,dM(t,x),$$

which is the multiple integral of order 1 defined by Itô [16]. This integral is centered, and for $g,h\in L^2([0,T]\times\mathbb{R},\mu)$,

$$E[M(h)M(g)]=\int_{[0,T]\times\mathbb{R}}h\,g\,d\mu.$$

We can write this integral as

$$M(h)=\sigma\int_0^T h(t,0)dW_t+\int_{[0,T]\times\mathbb{R}_0}h(t,x)x\,d\tilde{N}(t,x).$$

In this paper we also need to consider integrals with respect to the non compensated counting measure, defined by

$$\int_{[0,T]\times\mathbb{R}_0}h(t,x)x\,dN(t,x):=\int_{[0,T]\times\mathbb{R}_0}h(t,x)x\,d\tilde{N}(t,x)+\int_0^T\int_{\mathbb{R}_0}h(t,x)x\,dtd\nu(x),\tag{2.2}$$

when both integrals in the right hand side make sense.

Observe that if X has continuous trajectories, then M is nothing more than the independent Wiener measure, and if $\sigma=0$ and $\nu=\delta_1$ we obtain the standard random Poisson measure.

3 The derivative operators

3.1 Carlen-Pardoux derivative

The starting point of this paper is the following remark due to León and Tudor [20] about the Carlen–Pardoux derivative [8]: Let N be a Poisson process of parameter 1, and $\widetilde{N}_t=N_t-t$ be the compensated Poisson process. Denote by $T_1<\cdots<T_n\ldots$ the jump times of N. Let $h\in\mathcal{C}^1\big([0,T]\big)$. Then, the random variable

$$Y=\int_0^T h(s)\,d\widetilde{N}_s=\sum_{T_n\leq T}h(T_n)-\int_0^T h(s)\,ds$$

is in the domain of the Malliavin derivative in the Carlen-Pardoux sense and the derivative satisfies the following equality:

$$D_tY = \int_0^T h'(s)\big(\frac{s}{T} - 1_{(t,T]}(s)\big)\, dN_s.$$

In the next subsection, we extend formally this property to define a Malliavin derivative for a Lévy process.

3.2 The local Malliavin derivative

Henceforth, $\mathcal{S}$ denotes the family of all functionals of the form

$$f(M(h_1), \ldots, M(h_n)), \tag{3.1}$$

where f is in $C_b^\infty(\mathbb{R}^n)$ (f and all its partial derivatives are bounded), $h_1, \ldots, h_n \in L^2([0,T] \times \mathbb{R}, \mu)$ and, for all $x \neq 0$, $h_i(\cdot, x) \in C^1([0,T])$, $i \in \{1, \ldots, n\}$, and $\partial h_i \in L^2([0,T] \times \mathbb{R}, \mu)$ where ∂ means the partial derivative with respect to time. The set $\mathcal{S}$ is called the family of smooth random variables, and we will prove later that it is dense in $L^2(\Omega)$ (see Proposition 3.5).

We will also consider the family $\mathcal{K}$ of all bounded functions $k : [0,T] \times \mathbb{R}_0 \to \mathbb{R}$ such that they and their partial derivative with respect time are in $L^2([0,T] \times \mathbb{R}_0, \lambda \times \nu) \cap L^1([0,T] \times \mathbb{R}_0, \lambda \times \nu)$.

Now we introduce the main tool of this paper. Namely, the family of derivative operators for functionals of the Lévy process X.

Definition 3.1 *Given $k \in \mathcal{K}$, $F \in \mathcal{S}$ and $\Lambda \in \mathcal{B}(\mathbb{R})$, we define*

$$D_t^{\Lambda,k}F = \sum_{i=1}^n (\partial_i f)(M(h_1), \ldots, M(h_n)) D_t^{\Lambda,k} M(h_i), \quad t \in [0,T],$$

where

$$D_t^{\Lambda,k} M(h) = \mathbf{1}_\Lambda(0)\, \sigma h(t,0) + \int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\partial_s h(s,y)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))y\, dN(s,y). \tag{3.2}$$

We call $D_t^{\Lambda,k}F$ the local Malliavin derivative of F.

Remarks 3.2 1. The integral in the right-hand side of (3.2) is well-defined. Indeed, in agreement with (2.2),

$$\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\partial_s h(s,y)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))y\, d\tilde{N}(s,y)$$

is finite because k is bounded, and

$$\int_0^T \int_{\Lambda\cap\mathbb{R}_0} k(s,y)\partial_s h(s,y)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))y\, ds\nu(dy)$$

is finite by Cauchy-Schwarz inequality. Moreover, $D_t^{\Lambda,k}F \in L^2([0,T] \times \Omega, \lambda \otimes \mathbb{P})$. This explain why we need to introduce the family $\mathcal{K}$ in the definition of the derivative operator. Also in Section 5, we see that this family $\mathcal{K}$ allows to consider applications to stochastic differential equations.

2. The function $k(t,x)$ is an essential ingredient of the derivative, and it may change from one application to another. To shorten the notation, in general we will omit that k in the expression $D_t^{\Lambda,k}F$.

3. If $h, \partial h \in L^2([0,T]\times\mathbb{R}_0, \lambda\otimes\nu)$, then $h(s,y)/y$ and $\partial_s h(s,y)/y$ belong to $L^2([0,T]\times\mathbb{R}_0,\mu)$. Hence,

$$D_t^\Lambda M(h(s,y)/y) = \int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\partial_s h(s,y)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))dN(s,y),$$

   where the factor y has disappeared in the integral.

4. Observe that if $\Lambda = \{0\}$, then D^Λ is the Malliavin derivative with respect to the Brownian part of the Lévy process X.

5. If $\Lambda = \{x\}$ for some $x \neq 0$ with $\nu(\{x\}) \neq 0$, we obtain

$$D_t^{\{x\}} M(h) = x\int_0^T k(s,x)\partial_s h(s,x)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))dN_s^x,$$

   where N_s^x is the Poisson process on $[0,T]$ that counts the number of jumps of height x. Moreover, if the Lévy process is the standard Poisson process, $x = 1$, and if we take $k(t,x)$ independent of the time parameter, we obtain

$$D_t^{\{1\},k} M(h) = k(1)\int_0^T \partial_s h(s,1)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))dN_s^1.$$

   Like we commented in the previous subsection, that is exactly a constant times the derivative defined by Carlen and Pardoux in [8], as it is shown in León and Tudor [20].

6. The integration by parts formula allow us to write

$$\begin{aligned}
&\int_{\Lambda\cap\mathbb{R}_0}\int_0^T k(s,y)\partial_s h(s,y)y(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))ds\nu(dy) \\
&\qquad = \int_{\Lambda\cap\mathbb{R}_0} \big\{k(t,y)h(t,y) - \frac{1}{T}\int_0^T k(s,y)h(s,y)ds\big\} y\nu(dy) \\
&\qquad\quad - \int_0^T\int_{\Lambda\cap\mathbb{R}_0} h(s,y)\partial_s k(s,y)\left(\frac{s}{T} - \mathbf{1}_{[t,T]}(s)\right) y\nu(dy)ds.
\end{aligned}$$

   Thus,

$$\begin{aligned}
D_t^\Lambda M(h) &= \mathbf{1}_\Lambda(0)\,\sigma h(t,0) + \int_{[0,T]\times\Lambda\cap\mathbb{R}_0} k(s,y)\partial_s h(s,y)y(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))d\tilde{N}(s,y) \\
&\quad + \int_{\Lambda\cap\mathbb{R}_0} \big\{k(t,y)h(t,y) - \frac{1}{T}\int_0^T k(s,y)h(s,y)ds\big\} y\nu(dy) \\
&\quad - \int_0^T\int_{\Lambda\cap\mathbb{R}_0} h(s,y)\partial_s k(s,y)\left(\frac{s}{T} - \mathbf{1}_{[t,T]}(s)\right) y\nu(dy)ds.
\end{aligned}$$

7. If $0 \notin \Lambda$, then $D^\Lambda M(h)$ is orthogonal to the constant functions. In fact, the Fubini theorem (see Proposition 3.3) yields

$$\int_0^T \int_{[0,T]\times\Lambda} k(s,y)\partial_s h(s,y)y(\frac{s}{T} - \mathbf{1}_{[0,s]}(t))dN(s,y)dt$$
$$= \int_{[0,T]\times\Lambda} k(s,y)\partial_s h(s,y)y \int_0^T (\frac{s}{T} - \mathbf{1}_{[0,s]}(t))dtdN(s,y) = 0.$$

The next result is a Fubini type proposition. Its proof is straightforward and omitted.

Proposition 3.3 *Let $f \in L^2([0,T]^2 \times \mathbb{R}, \lambda^{\otimes 2} \otimes \mu)$. Then the following integrals are well defined and*

$$\int_0^T M(f(t,\cdot))\,dt = M(\int_0^T f(t,\cdot)\,dt), \text{ a.s.} \tag{3.3}$$

In the Brownian Malliavin calculus if F is a smooth random variable such that $F = 0$, it follows that $DF = 0$. However, for Lévy processes matters are different. For example, if X is a standard Poisson process and we take $h = 1$, then $F = f(M(h)) = f(N_T - T)$, and $F = 0$ only implies that f is zero on the set $\mathbb{N} - T$. In the next proposition we show the corresponding property.

Proposition 3.4 *Let $F \in \mathcal{S}$ such that $F = 0$ a.s. Then $D^\Lambda F = 0,\ \lambda \otimes \mathbb{P} -$ a.e.*

Proof. Let $F = f(M(h_1),\ldots,M(h_n))$. To consider the most general case, assume that $0 \in \Lambda$. By definition,

$$D_t^\Lambda F = \underbrace{\sum_i (\partial_i f)(M(h_1),\ldots,M(h_n))\sigma h_i(t,0)}_{(*)}$$
$$+ \underbrace{\sum_{i=1}^n (\partial_i f)(M(h_1),\ldots,M(h_n)) \int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\partial_s h_i(s,y)(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))y\,dN(s,y)}_{(**)}.$$

The term (*) is a Malliavin derivative with respect to the Brownian part (see Solé *et al.* [26]). From the independence between the Brownian part and the jumps part of X, by fixing the random variables corresponding to the jumps part, we have that

$$M(h_j) = \sigma W(h_j) + C_j,$$

where $W(h) = \int_0^T h(t,0)\,dW_t$ and C_j is a constant; changing the function f we can take $C_1 = \cdots = C_n = 0$. First assume that the Gaussian vector $(W(h_1),\ldots,W(h_n))$ has a regular covariance matrix. Then it has density, and

$$f(W(h_1)\ldots,W(h_n)) = 0, \text{ a.s.}$$

implies

$$f(x_1,\ldots,x_n) = 0,\ \lambda^n - \text{a.e.}$$

It follows that all derivatives of f are zero.

If the covariance matrix of $(W(h_1),\dots,W(h_n))$ is singular, denote by k the rank of that matrix, and (reordering) assume that $(W(h_1),\dots,W(h_k))$ has a density. Then there are numbers μ_i^j, $i = 1,\dots,n$, $j = k+1,\dots,n$ such that

$$W(h_j) = \sum_{i=1}^{k} \mu_i^j W(h_i),\ j = k+1,\dots,n.$$

Given the linearity of Itô-Wiener integral, it follows that λ–a.s.,

$$h_j = \sum_{i=1}^{k} \mu_i^j h_i,\ j = k+1,\dots,n.$$

Consider the vectorial function

$$\boldsymbol{\Phi} = (\Phi_1,\dots,\Phi_n) : \mathbb{R}^k \to \mathbb{R}^n$$

given by

$$\Phi_j(x_1,\dots,x_k) = x_j,\ \text{for}\ j = 1,\dots,k,$$

and

$$\Phi_j(x_1,\dots,x_k) = \sum_{i=1}^{k} \mu_i^j x_i,\ j = k+1,\dots,n.$$

Define $g := f \circ \boldsymbol{\Phi}$. Then

$$f(W(h_1)\dots,W(h_n)) = g(W(h_1)\dots,W(h_k)).$$

Since $\big(W(h_1)\dots,W(h_k)\big)$ has a density, we have that

$$\frac{\partial g}{\partial x_j} = 0,\ j = 1,\dots,k.$$

By the chain rule,

$$J_g = J_f\, J_{\boldsymbol{\Phi}},$$

where J_m denotes the Jacobian matrix of a generic function m. Writing $\boldsymbol{h} = (h_1,\dots,h_k)^T$, we have

$$0 = J_g \boldsymbol{h}(t) = J_f J_{\boldsymbol{\Phi}} \boldsymbol{h}(t) = \sum_{j=1}^{n} \frac{\partial f}{\partial x_j}\, h_j(t),$$

and hence (*) is zero.

The term (**) is treated as follows. Again by the independence between the Brownian and the jumps part, we can assume $\sigma = 0$. For the sake of simplicity we consider the case $n = 2$.

Consider the set

$$\Lambda_m = \Lambda \cap \{x \in \mathbb{R} : 1/m < |x| < m\},$$

and denote by N^{Λ_m} the restriction of the counting measure of jumps N to the jumps with size in Λ_m. That is, for $B \in \mathcal{B}\big((0,\infty)\times\mathbb{R}_0\big)$,

$$N^{\Lambda_m}(B) = \#\{t :\ (t,\Delta X_t) \in B \text{ and } \Delta X_t \in \Lambda_m\},$$

and

$$N_T^m = N^{\Lambda_m}([0,T]\times\mathbb{R}_0) < \infty, \text{ a.s.},$$

due to $\nu(\Lambda_m) < \infty$. The jumps of X in $[0,T]$ with height in Λ_m can be ordered, $T_1 < T_2 < \cdots < T_{N_T^m}$ Given $N_T^m = k$, the vector $(T_1,\ldots,T_k)$ has a Dirichlet distribution on $[0,T]$, independent of the height of the jumps. The jump size has a distribution given by $P\{\Delta X_{T_j} \in A\} = \nu(A)/\nu(\Lambda_m)$, $A \in \mathcal{B}(\Lambda_m)$.

In a similar way, define $N^{\mathbb{R}_0-\Lambda_m}$. Since N^{Λ_m} and $N^{\mathbb{R}_0-\Lambda_m}$ are independent, denoting by $\mathbb{E}_{\Lambda_m}$ and $\mathbb{E}_{\mathbb{R}_0-\Lambda_m}$ the expectation with respect to each jump counting measure, and as $F=0$ implies $F^2 = f^2(M(h_1),\ldots,M(h_n)) = 0$, we have that

$$0 = \mathbb{E}\big[F^2\big] = \mathbb{E}_{\mathbb{R}_0-\Lambda_m}\mathbb{E}_{\Lambda_m}\big[F^2\big] = \mathbb{E}_{\mathbb{R}_0-\Lambda_m}\Big[\sum_{k\geq 0}\mathbb{P}\{N_T^m = k\}\mathbb{E}\big[F^2\,|\,N_T^m = k\big]\Big].$$

Due to the fact that $F^2 \geq 0$, for each k,

$$\begin{aligned}0 = \mathbb{E}_{\mathbb{R}_0-\Lambda_m}\Big[\int_{S_k(\Lambda_m)} f^2\Big(&\sum_{j=1}^k h_1(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_1(s,y)y\,d\nu(y)ds + M\big(h_1\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big),\\ &\sum_{j=1}^k h_2(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_2(s,y)y\,d\nu(y)ds + M\big(h_2\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big)\Big)\,ds_1\cdots ds_k d\nu(y_1)\cdots d\nu(y_k)\Big],\end{aligned}$$

where $S_k(\Lambda_m)$ the *simplex*

$$\big\{(t_1,x_1;\ldots;t_k,x_k) \in ([0,T]\times\Lambda_m)^k:\ t_1 < \cdots < t_k\big\}.$$

Therefore, using again that f^2 is non negative, $P_{\mathbb{R}_0-\Lambda_m}$–a.s., the function from $S_k(\Lambda_m)$ into $\mathbb{R}$,

$$\begin{aligned}(s_1,y_1;\ldots;s_k,y_k) \mapsto f\Big(&\sum_{j=1}^k h_1(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_1(s,y)y\,d\nu(y)ds + M\big(h_1\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big),\\ &\sum_{j=1}^k h_2(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_2(s,y)y\,d\nu(y)ds + M\big(h_2\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big)\Big)\end{aligned}$$

is zero a.e. $ds_1\ldots ds_k\,\nu(dy_1)\ldots\nu(dy_k)$. Hence, by the continuity of both f and $h_i(\cdot,y)$, it is deduced that, fixed $(y_1,\ldots,y_k)$, ν^k–a.e., the derivatives with respect to the time variables s are also zero. and then, for every $\ell = 1,\ldots,k$

$$\begin{aligned}\sum_{i=1}^2 \partial_i f\Big(&\sum_{j=1}^k h_1(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_1(s,y)y\,d\nu(y) + M\big(h_1\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big),\\ &\sum_{j=1}^k h_2(s_j,y_j)y_j - \int_0^T\int_{\Lambda_m} h_2(s,y)y\,d\nu(y) + M\big(h_2\mathbf{1}_{\mathbb{R}_0-\Lambda_m}\big)\Big)\partial_s h_i(s_\ell,y_\ell)y_\ell = 0,\end{aligned}$$

a.e. $ds_1\ldots ds_k\,\nu(dy_1)\ldots\nu(dy_k)$. Multiplying this quantity by $k(s_\ell,y_\ell)\big(\frac{s_\ell}{T} - \mathbf{1}_{(t,T]}(s_\ell)\big)$ and adding with respect to ℓ, we arrive to

$$\sum_{i=1}^2 \partial_i f\big(M(h_1),M(h_2)\big)\int_{[0,T]\times\Lambda_m} k(s,y)\partial_s h_i(s,y)y\big(\frac{s}{T} - \mathbf{1}_{(t,T]}(s)\big)\,dN(s,y) = 0, \text{ a.s.}$$

Now, thanks to Remark 3.2, we can apply the dominated convergence Theorem to pass to the limit that expression when $m\to\infty$, and we obtain that (**) is 0. □

Proposition 3.5 *$\mathcal{S}$ is dense in $L^2(\Omega)$.*

Proof. Since the finite linear combinations of multiple integrals are dense in $L^2(\Omega)$, it is enough to prove that the random variables of the form $M(E_1)\cdots M(E_n)$ for $E_1,\dots,E_n \in \mathcal{B}\big([0,T]\times\mathbb{R}\big)$ disjoint sets, with finite μ measure, can be approximated by elements of $\mathcal{S}$. We proceed in four steps.

1. It suffices to prove the approximation for disjoint closed sets. This is proved in the following way. The measure μ is σ–finite in $\mathcal{B}\big([0,T]\times\mathbb{R}\big)$, and hence it is regular in the sense that for every $A\in\mathcal{B}\big([0,T]\times\mathbb{R}\big)$,

$$\mu(A)=\sup\{\mu(C):\, C\subset A,\, C \text{ closed}\}=\inf\{\mu(O):\, A\subset O,\, O \text{ open}\}.$$

So, given $E_1,\dots,E_n \in \mathcal{B}\big([0,T]\times\mathbb{R}\big)$, for every $\varepsilon>0$, there are closed sets $C_1,\dots,C_n$, with $C_j\subset E_j$ and

$$\mu(E_j-C_j)<\varepsilon,\ j=1,\dots,n.$$

From the fact that $E_1,\dots,E_n$ are disjoints and $C_j\subset E_j$, it follows that

$$E\Big[\big(M(E_1)\cdots M(E_n)-M(C_1)\cdots M(C_n)\big)^2\Big]\le n\max_{j=1,\dots,n}\{\mu(E_j)\}^{n-1}\max_{j=1,\dots,n}\{\mu(E_j-C_j)\},$$

and the claim of this step follows.

2. We can assume that $E_1,\dots,E_n$ are disjoint open sets: From previous step, we can take $E_1,\dots,E_n$ as disjoint closed sets. By the separability properties of $\mathbb{R}^n$, they can be separated by open sets, that is, there are disjoint open sets $O_1,\dots,O_n$ such that $E_j\subset O_j$. By the regularity of μ, and the same reasoning as before, the property is proved.

3. Let $E_1,\dots,E_n$ be disjoint open sets. Since every open set on $[0,T]\times\mathbb{R}$ is a countable union of open rectangles, we can assume that each E_j is a finite union of open rectangles (with finite μ-measure) of the form $(t',t'')\times(x',x'')$, or $[0,t)\times(x',x'')$ or $(t,T]\times(x',x'')$; moreover, we can reduce slightly the size of all the t-intervals in such a way that the measures $\mu(E_j),\ j=1,\dots,n$, do not change significantly: specifically, given $\varepsilon>0$, there is $\delta>0$ such that if we denote E_j^δ the union with rectangles $(t'+\delta,t''-\delta)\times(x',x'')$ instead of the corresponding $(t',t'')\times(x',x'')$ of E_j, and similarly for the other types of rectangles, then

$$\mu(E_j-E_j^\delta)\le\varepsilon,\ j=1,\dots,n.$$

Let $m_r\in\mathcal{C}_0^\infty(\mathbb{R}),\ \ r>0$ a family of mollifiers (see, for example, Adams [1, Page 29]). For a rectangle $(t'+\delta,t''-\delta)\times(x',x'')$ write

$$f_r(t,x)=\Big(m_r*\mathbf{1}_{(t'+\delta,t''-\delta)}\Big)(t)\,\mathbf{1}_{(x',x'')}(x),$$

where the symbol $*$ means convolution. Collecting all the f_r corresponding to the same E_j^δ, and using the fact that the support are included in E_j for $r<\delta$, we obtain that a random variable of the form $M(g_1)\cdots M(g_n)$, with $g_1,\dots g_n$ with the required properties, and moreover, with disjoint support, approximates the product $M(E_1)\cdots M(E_n)$ in $L^2(\Omega)$.

4. Finally, use an approximation by $\mathcal{C}_0^\infty(\mathbb{R}^n)$ functions to the product function to get the Proposition. □

Remark 3.6 In the preceding proof we have proved that instead of $\mathcal{S}$ we can use the sum of the smooth random variables of the form $f(M(h_1),\dots,M(h_n))$ with $h_1,\dots,h_n$ with disjoint support to define the local Malliavin derivative.

Next proposition gives a Malliavin integration by parts formula:

Proposition 3.7 *Let $F \in \mathcal{S}$, and g be a measurable and bounded function on $[0,T]$. Then*

$$\mathbb{E}\Big[\int_0^T (D_t^\Lambda F)g(t)dt\Big] \tag{3.4}$$

$$= \mathbb{E}\Big[F\Big(\mathbf{1}_\Lambda(0)\sigma\int_0^T g(s)dW_s + \int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\big(g(s) - \frac{1}{T}\int_0^T g(t)dt\big)k(s,y)dN(s,y)\Big)\Big] \tag{3.5}$$

$$- \mathbb{E}\Big[F\Big(\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\partial_s k(s,y)\big[\int_0^T g(t)(\frac{s}{T} - \mathbf{1}_{[0,s]}(t))dt\big]dN(s,y)\Big)\Big]. \tag{3.6}$$

Proof:

First we consider the case $F = f(M(h))$, where f and h are as in (3.1) and assume $\nu(\Lambda) < \infty$. Then,

$$\begin{aligned}
&\mathbb{E}\left[\int_0^T (D_t^\Lambda f(M(h)))g(t)dt\right] \\
&= \mathbf{1}_\Lambda(0)\mathbb{E}\left[\sigma\int_0^T f'(M(h))h(t,0)g(t)dt\right] \\
&\quad + \mathbb{E}\left[\int_0^T f'(M(h))\left(\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\partial_s h(s,y)y(\frac{s}{T} - \mathbf{1}_{[t,T]}(s))dN(s,y)\right)g(t)dt\right]
\end{aligned} \tag{3.7}$$

From the integration by parts formula in the Wiener case,

$$\mathbb{E}\left[\int_0^T D_t^{\{0\}} f(M(h))g(t)dt\right] = \mathbb{E}\left[f(M(h))M(g(t)\otimes\mathbf{1}_{\{0\}}(x))\right]$$

and the first term of the right hand side of (3.7) is equal to

$$\mathbf{1}_\Lambda(0)\mathbb{E}[f(M(h))M(g\otimes\mathbf{1}_{\Lambda\cap\{0\}})].$$

Now, using Fubini theorem, the second term can be written as

$$\mathbb{E}\Big[f'(M(h))\Big(\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\Big\{k(s,y)\partial_s h(s,y)y\big(\frac{s}{T}\int_0^T g(t)dt - \int_0^s g(t)dt\big)\Big\}dN(s,y)\Big)\Big],$$

which, by the independence of W, N^Λ and $N^{\mathbb{R}_0-\Lambda}$ (using the same notations as in the previous proof), and the symmetry of the function $\sum_{j=1}^n h(s_j,y_j)y_j$, is equal to

$$\begin{aligned}
&\sum_{n=1}^\infty \mathbb{P}\{N_T^{\Lambda\cap\mathbb{R}_0} = n\}\frac{1}{\nu(\Lambda)^n T^n} \\
&\quad\times\int_{[0,T]^n}\int_{(\Lambda\cap\mathbb{R}_0)^n}\Big\{\mathbb{E}_{W,N^{\mathbb{R}_0-\Lambda}}\Big[f'(\sum_{j=1}^n h(s_j,y_j)y_j - \int_0^T\int_\Lambda h(r,z)z\,drd\nu(z) \\
&\qquad\qquad + \mathbf{1}_\Lambda(0)M(h\mathbf{1}_{\{0\}}) + M(h\mathbf{1}_{\mathbb{R}_0-\Lambda}))\Big] \\
&\quad\times\sum_{j=1}^n k(s_j,y_j)\partial_s h(s_j,y_j)y_j\Big(\frac{s_j}{T}\int_0^T g(t)dt - \int_0^{s_j} g(t)dt\Big)\Big\}\,ds_1\cdots ds_n d\nu(y_1)\cdots d\nu(y_n).
\end{aligned}$$

Integrating by parts each integral

$$\int_0^T \mathbb{E}_{W,N^{\mathbb{R}_0-\Lambda}}\Big[f'(\sum_{j=1}^n h(s_j,y_j)y_j - \int_0^T\int_\Lambda h(r,z)z\,drd\nu(z)+\cdots)\Big] \\ \times k(s_i,y_i)\partial_s h(s_i,y_i)y_i\Big(\frac{s_i}{T}\int_0^T g(t)dt - \int_0^{s_i} g(t)dt\Big)ds_i,$$

and reordering terms, we arrive to

$$\mathbb{E}\Big[f(M(h))\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)} k(s,y)\Big(g(s)-\frac{1}{T}\int_0^T g(t)dt\Big)dN(s,y)\Big] \\ -\mathbb{E}\Big[f(M(h))\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\partial_s k(s,y)\Big(\int_0^T g(t)(\frac{s}{T}-\mathbf{1}_{[0,s]}(t))dt\Big)dN(s,y)\Big],$$

which implies that the result holds in the case that $\nu(\Lambda)<\infty$.

To deal with the case $\nu(\Lambda)=\infty$, let $\Lambda_n=\{0\}\cup\{\frac{1}{n}<|x|<n\}$. Then, changing Λ by $\Lambda_n\cap\Lambda$, and letting $n\to\infty$, we obtain the result.

Now, consider $F=f(M(h_1),\ldots,M(h_n))$, with $h_1,\ldots,h_n$ with disjoint support. Since the random variables $M(h_1),\ldots,M(h_n)$ are independent, we can compute the expectations in (3.4) iteratively, and the result follows. Finally, by linearity, the equality (3.4) is extended to a sum of smooth random variables of the form $f(M(h_1),\ldots,M(h_n))$, with $h_1,\ldots,h_n$ with disjoint support, which generate $\mathcal{S}$, in agreement with Remark 3.6. □

The following result is a consequence of Proposition 3.7 applied to FG and of the fact that D^Λ is a true derivative operator, which follows from Definition 3.1.

Corollary 3.8 *Let g measurable and bounded, and $F,G\in\mathcal{S}$. Then*

$$\mathbb{E}\left(G\int_0^T(D_t^\Lambda F)g(t)dt + F\int_0^T(D_t^\Lambda G)g(t)dt\right) \\ =\mathbb{E}\left[FG\left(\mathbf{1}_\Lambda(0)\int_0^T\sigma g(s)dW_s + \int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\left(g(s)-\frac{1}{T}\int_0^T g(t)dt\right)k(s,y)dN(s,y)\right.\right. \\ \left.\left.-\int_{[0,T]\times(\Lambda\cap\mathbb{R}_0)}\partial_s k(s,y)\left(\int_0^T g(t)(\frac{s}{T}-\mathbf{1}_{[0,s]}(t))dt\right)dN(s,y)\right)\right].$$

Now we can proceed as in Nualart [22] (pag. 26) to show the following result, taking into account that the bounded functions are dense in $L^2([0,T])$.

Corollary 3.9 *The operator D^Λ, from $L^2(\Omega)$ into $L^2([0,T]\times\Omega)$, is unbounded, densely defined and closable.*

In particular, we have that the operator D^Λ has a closed extension, which is also written by D^Λ. The domain of this operator is denoted by $\mathrm{Dom}\,D^\Lambda$. Note that F belongs to $\mathrm{Dom}\,D^\Lambda$ if and only if there is a sequence $\{F_n : n\geq 1\}\subset\mathcal{S}$ such that F_n converges to F in $L^2(\Omega)$, when $n\to\infty$, and $\{D^\Lambda F_n : n\geq 1\}$ converges to some process Y in $L^2([0,T]\times\Omega)$. In this case, we have that $D^\Lambda F=Y$.

The following rules of derivation are well known in Brownian Malliavin calculus; they can be translated to our context with the same proof.

Proposition 3.10 **1.** *(Chain rule). Let $f : \mathbb{R}^n \to \mathbb{R}$ be a continuously differentiable function with bounded partial derivatives, and let $\boldsymbol{F} = (F_1, \dots, F_n)$ a random vector such that $F_j \in \operatorname{Dom} D^\Lambda$, for $j = 1, \dots, n$. Then $f(\boldsymbol{F}) \in \operatorname{Dom} D^\Lambda$ and*

$$D^\Lambda\big(f(\boldsymbol{F})\big) = \sum_{j=1}^n \partial_j f(\boldsymbol{F}) D^\Lambda F_j.$$

2. *Let $F, G \in \operatorname{Dom} D^\Lambda$ such that $G\, D^\Lambda F$ and $F\, D^\Lambda G$ belong to $L^2([0,T]\times\Omega)$. Then $F\,G \in \operatorname{Dom} D^\Lambda$ and*

$$D^\Lambda(F\,G) = G\, D^\Lambda F + F\, D^\Lambda G.$$

3.3 Absolute continuity via the operators D^Λ

The chain rule allows to prove a criterion for the absolutely continuity of a functional F in a similar way that in the Brownian case. Specifically, the result obtained by Carlen and Pardoux [8, Theorem 4.1] for the Poisson process can be extended to the Lévy case; the proof is the same as Carlen and Pardoux, and omitted.

Proposition 3.11 *Let $\Lambda \in \mathcal{B}(\mathbb{R})$, $k \in \mathcal{K}$ and $F \in \operatorname{Dom} D^\Lambda$ such that*

$$\int_0^T (D_t^\Lambda F)^2 dt > 0$$

a.s. on a measurable set $A \in \mathcal{F}$. Then, $\mathbb{P} \circ F^{-1}$ is absolutely continuous on A, that is, $\lambda(B) = 0$ implies $\mathbb{P}(\{F \in B\} \cap A) = 0$.

It is worth to remark that the criterion is true for every set Λ and weight function $k \in \mathcal{K}$. This is very interesting for applications because we can choose an appropriate Λ and k depending of F. It will be used in Section 5 to see that the solutions of some stochastic differential equations have a density on a measurable set.

4 Multiple integrals over sets with finite Lévy measure and its derivatives

In this section we study multiple integrals defined on a set with finite Lévy measure. Such integrals can be computed pathwise, and its properties can be proved by combinatorial methods.

Let $\Theta \in \mathcal{B}(\mathbb{R})$ be a bounded set such that $0 \notin \overline{\Theta}$ (closure of Θ); then $\nu(\Theta) < \infty$. As we have done before, consider

$$N^\Theta(B) = \#\{t : \ (t, \Delta X_t) \in B \text{ and } \Delta X_t \in \Theta\}, \text{ for } B \in \mathcal{B}\big((0,\infty) \times \mathbb{R}_0\big),$$

that is a Poisson random measure with intensity $\lambda \otimes \nu_\Theta$, where $\nu_\Theta(C) = \nu(\Theta \cap C)$, for $C \in \mathcal{B}(\mathbb{R})$. Write

$$N_T^\Theta = N^\Theta\big([0,T] \times \Theta\big) < \infty, \text{ a.s.}$$

Then we can order the jumps in the interval $[0,T]$, say $T_1 < \cdots < T_{N_T^\Theta}$, and write

$$N^\Theta = \sum_{j=1}^{N_T^\Theta} \delta_{(T_j, \Delta X_{T_j})}.$$

Let $\phi : [0,T] \times \mathbb{R}_0 \to \mathbb{R}$ be a measurable function (indeed, it is only needed that ϕ is defined in $[0,T] \times \Theta$) integrable with respect to N^Θ and consider

$$J_1^\Theta(\phi) := \int_{[0,T]\times\mathbb{R}_0} \phi(t,x)\, dN^\Theta(t,x).$$

This integral has a compound Poisson distribution with characteristic function (see, for example, Sato [27, Proposition 19.5])

$$\mathbb{E}\big[e^{iuJ_1^\Theta(\phi)}\big] = \exp\Big\{ \int_{[0,T]\times\Theta} \big(e^{iu\phi(t,x)} - 1\big)\, dtd\nu(x)\Big\}.$$

Thus, if $\phi \in L^p([0,T]\times\mathbb{R}_0, \lambda\otimes\nu_\Theta)$, then $J_1^\Theta(\phi) \in L^p(\Omega)$, for $p \geq 1$.

Also multiple integrals can be considered. Recall the notations introduced in the proof of Proposition 3.4: let $S_n(\Theta)$ be the *simplex*

$$\big\{(t_1,x_1;\ldots;t_n,x_n) \in ([0,T]\times\Theta)^n :\ 0 \leq t_1 < \cdots < t_n \leq T\big\}.$$

For $\phi : S_n(\Theta) \to \mathbb{R}$, define

$$\begin{aligned} J_n^\Theta(\phi) &= \int\cdots\int_{S_n(\Theta)} \phi(t_1,x_1;\ldots;t_n,x_n)\, dN^\Theta(t_1,x_1)\cdots dN^\Theta(t_n,x_n) \\ &= \sum_{1\leq i_1<\cdots<i_n\leq N_T^\Theta} \phi(T_{i_1},\Delta X_{T_{i_1}};\ldots;T_{i_n},\Delta X_{T_{i_n}})\, \mathbf{1}_{\{N_T^\Theta\geq n\}}. \end{aligned}$$

Proposition 4.1 *For every $p\geq 1$, if $\phi\in L^p\big(S_n(\Theta),(\lambda\otimes\nu_\Theta)^n\big)$, then $J_n^\Theta(\phi)\in L^p(\Omega)$, and*

$$\mathbb{E}\big[\big|J_n^\Theta(\phi)\big|^p\big] \leq C_{p,n}\int_{S_n(\Theta)} |\phi(t_1,x_1;\ldots;t_n,x_n)|^p\, dt_1\cdots dt_n d\nu(x_1)\cdots d\nu(x_n), \tag{4.1}$$

where the constant $C_{n,p}$ depends only on n and p.

Proof. The main tool of the proof is that the jump times of a compound Poisson process follow a Dirichlet distribution, independent of the size of jumps, which are also independent and each of one has law $P\{\Delta X_{T_j}\in A\} = \nu(A)/\nu(\Theta)$, for $A\in\mathcal{B}(\Theta)$. To simplify the notations, we assume $T=1$. By Jensen inequality,

$$|J_n^\Theta(\phi)|^p \leq \Big(\Big(N^{(\Theta)}\otimes\cdots\otimes N^\Theta)(S_n(\Theta))\Big)^{p-1}\int_{S_n(\Theta)} |\phi(t_1,x_1;\ldots;t_n,x_n)|^p\, dN^\Theta(t_1,x_1)\cdots dN^\Theta(t_n,x_n).$$

Hence,

$$\begin{aligned} \mathbb{E}\big[\big|J_n^\Theta(\phi)\big|^p\big] &= \sum_{k\geq n}\mathbb{E}\Big[\big|J_n^\Theta(\phi)\big|^p \,|\, N_1^\Theta = k\Big]\,\mathbb{P}\{N_1^\Theta = k\} \\ &\leq e^{-\nu(\Theta)}\sum_{k\geq n}\frac{\nu(\Theta)^k}{k!}\,k^{p-1}\frac{1}{\nu(\Theta)^n}\,\mathbb{E}\Big[\sum_{1\leq i_1<\cdots<i_n\leq k}\big|\phi(T_{i_1},\Delta X_{T_{i_1}};\ldots;T_{i_n},\Delta X_{T_{i_n}})\big|^p\Big] \\ &= e^{-\nu(\Theta)}\sum_{k\geq n}\frac{\nu(\Theta)^k}{k!}\,k^{p-1}\frac{1}{\nu(\Theta)^n}\sum_{1\leq i_1<\cdots<i_n\leq k}\int_{\Theta^n}\mathbb{E}\Big[\big|\phi(U_{(i_1)},x_1;\ldots;U_{(i_n)},x_n)\big|^p\Big]d\nu(x_1)\ldots d\nu(x_n), \end{aligned}$$

where $(U_{(i_1)}, \dots, U_{(i_n)})$ is the ordered $(i_1, \dots, i_n)$–statistic from a sample $U_1, \dots, U_k$ of a uniform law in $(0, 1)$. Extend ϕ to the whole $(0, 1)^n$ by symmetry, and denote by $\mathcal{C}_n^k$ the family of all subsets of $\{1, \dots, k\}$ of size n. Fixed $x_1, \dots, x_n \in \Theta$,

$$\begin{aligned}
\mathbb{E}\Big[\sum_{1\le i_1<\cdots<i_n\le k} \big|\phi(U_{(i_1)}, x_1; \dots; U_{(i_n)}, x_n)\big|^p\Big] &= \mathbb{E}\Big[\sum_{\{i_1,\dots,i_n\}\in\mathcal{C}_n^k} \big|\phi(U_{i_1}, x_1; \dots; U_{i_n}, x_n)\big|^p\Big] \\
&= \binom{k}{n} \int_{(0,1)^n} \big|\phi(t_1, x_1, \dots, t_n, x_n)\big|^p \, dt_1 \cdots dt_n \\
&= \frac{k!}{(k-n)!} \int_{0<t_1\cdots<t_n<1} \big|\phi(t_1, x_1, \dots, t_n, x_n)\big|^p \, dt_1 \cdots dt_n.
\end{aligned}$$

Finally,

$$\begin{aligned}
\mathbb{E}\big[\big|J_n^\Theta(\phi)\big|^p\big] \le \int_{0<t_1\cdots<t_n<1} \int_{\Theta^n} \big|\phi(t_1, x_1, \dots, t_n, x_n)\big|^p \, dt_1 \cdots dt_n d\nu(x_1) \dots d\nu(x_n) \\
\times e^{-\nu(\Theta)} \sum_{k\ge n} \frac{\nu(\Theta)^{k-n}}{(k-n)!} k^{p-1},
\end{aligned}$$

and the result follows. □

Corollary 4.2 *Let $\phi_k, \phi \in L^p\big(S_n(\Theta), (\lambda \otimes \nu_\Theta)^n\big)$, such that*

$$\lim_k \phi_k(t_1, x_1; \dots; t_n, x_n) = \phi(t_1, x_1; \dots; t_n, x_n), \text{ uniformly}.$$

Then

$$\lim_k J_n(\phi_k) = J_n(\phi), \text{ in } L^p(\Omega).$$

A crucial result in multiple integration theory is the product formula (see Lee and Shih [19] and Tudor [28]). For this particular case, the formula is simpler and the proof straightforward:

Proposition 4.3 *Consider $\phi_n : S_n(\Theta) \to \mathbb{R}$ and $\phi_1 : [0, T] \times \mathbb{R}_0 \to \mathbb{R}$. Then,*

$$J_n^\Theta(\phi_n)\, J_1^\Theta(\phi_1) = J_{n+1}^\Theta(\phi_n \widetilde{\otimes} \phi_1) + J_n^\Theta(\phi_n * \phi_1), \tag{4.2}$$

where

$$\phi_n \widetilde{\otimes} \phi_1(t_1, x_1; \dots; t_{n+1}, x_{n+1}) = \sum_{j=1}^{n+1} \phi_n(t_1, x_1; \dots; \widehat{t_j, x_j}; \dots; t_{n+1}, x_{n+1}) \phi_1(t_j, x_j),$$

where $\widehat{t_j, x_j}$ means that this pair is missing, and

$$\phi_n * \phi_1(t_1, x_1; \dots; t_n, x_n) = \sum_{j=1}^{n} \phi_n(t_1, x_1; \dots; t_n, x_n) \phi_1(t_j, x_j).$$

Remark 4.4 Assume that ϕ is defined in all $([0,T]\times\Theta)^n$. The function $\phi_n\widetilde{\otimes}\phi_1$ is not the standard symmetrization of $\phi_n\otimes\phi_1$, which is given by

$$\widetilde{\phi_n\otimes\phi_1}(t_1,x_1;\dots;t_{n+1},x_{n+1})=\frac{1}{(n+1)!}\sum_\sigma \phi_n(t_{\sigma_1},x_{\sigma_1};\dots;t_{\sigma_n},x_{\sigma_n})\phi_1(t_{\sigma_{n+1}},x_{\sigma_{n+1}}),$$

where the sum ranges over the set of permutations of $n+1$ elements. We have the relation

$$\widetilde{\phi_n\otimes\phi_1}=\frac{1}{n+1}\,\widetilde{\phi}_n\widetilde{\otimes}\phi_1,\ \text{where } \widetilde{\phi}_n \text{ is the symmetrization of } \phi_n.$$

Lemma 4.5 *Let $\Theta\in\mathcal{B}(\mathbb{R})$ be a bounded set such that $0\notin\overline{\Theta}$, and $\Lambda\in\mathcal{B}(\mathbb{R})$ with $\Theta\subset\Lambda$. Consider $f\in\mathcal{C}_p^\infty(\mathbb{R})$. Then $f(N_T^\Theta)\in\operatorname{Dom}D^\Lambda$ and*

$$D^\Lambda f(N_T^\Theta)=0.$$

Proof. Write

$$\begin{aligned}N_T^\Theta&=\int_{[0,T]\times\mathbb{R}_0}dN^\Theta(t,x)=\int_{[0,T]\times\mathbb{R}_0}\frac{1}{x}\mathbf{1}_\Theta(x)x\,dN(t,x)\\&=\int_{[0,T]\times\mathbb{R}_0}\frac{1}{x}\mathbf{1}_\Theta(x)x\,d\widetilde{N}(t,x)+\int_0^T\int_\Theta dtd\nu(x).\end{aligned}$$

Hence, $N_T^\Theta\in\mathcal{S}$, and $D^\Lambda N_T^\Theta=0$. Thus also $f(N_T^\Theta)\in\mathcal{S}$ and $D^\Lambda f(N_T^\Theta)=0$. ☐

Given a set $A\subset\mathbb{N}$, the random variable $\mathbf{1}_{\{N_T^\Theta\in A\}}$ can be conveniently approximated by functionals $f(N_t^\Theta)\in\operatorname{Dom}D^\Lambda$. From the closability of D^Λ we get the following corollary:

Corollary 4.6 *For every set $A\subset\mathbb{N}$, $\mathbf{1}_{\{N_T^\Theta\in A\}}\in\operatorname{Dom}D^\Lambda$ and $D^\Lambda\mathbf{1}_{\{N_T^\Theta\in A\}}=0$.*

Theorem 4.7 *Let Θ and Λ as in Lemma 4.5, and $\phi\in L^2\big(S_n(\Theta),(\lambda\otimes\nu)^n\big)$ such that for every $(x_1,\dots,x_n)\in\Theta^n$, $\phi(\cdot,x_1;\dots;\cdot,x_n)\in\mathcal{C}^\infty\big(\overline{S_n}\big)$, where $\overline{S_n}=\{0\le t_1\le\cdots\le t_n\le T\}$. Then $J_n^\Theta(\phi)\in\operatorname{Dom}D^\Lambda$ and*

$$D_t^\Lambda J_n^\Theta(\phi)=\sum_{j=1}^n J_n^\Theta\Big(k(s_j,x_j)\partial_{s_j}\phi(s_1,x_1;\dots;s_n,x_n)\big(\frac{s_j}{T}-\mathbf{1}_{(t,T]}(s_j)\big)\Big).\tag{4.3}$$

Proof. We proceed by induction. For $n=1$, as in the previous lemma, write

$$\begin{aligned}J_1^\Lambda(\phi)&=\int_{[0,T]\times\mathbb{R}_0}\phi(s,x)\,dN^\Theta(s,x)\\&=M(\mathbf{1}_\Theta(x)\phi(s,x)/x)+\int_0^T\int_\Theta\phi(s,x)\,dtd\nu(x).\end{aligned}$$

Its derivative is

$$\int_{[0,T]\times\mathbb{R}_0}\mathbf{1}_\Theta(x)k(s,x)\partial_s\phi(s,x)\big(\frac{s}{T}-\mathbf{1}_{(t,T]}(s)\big)\,dN(s,x),$$

which coincides with (4.3).

Now, consider $\phi_n \in \mathcal{C}_0^\infty\big(S_n(\Theta)\big)$ and $\phi_1 \in \mathcal{C}_0^\infty\big((0,T)\times R_0\big)$. Then the result is obtained using the product formula (4.2)

$$J^\Theta_{n+1}(\phi_n\widetilde{\otimes}\phi_1) = J^\Theta_n(\phi_n)\, J^\Theta_1(\phi_1) - J^\Theta_n(\phi_n * \phi_1), \tag{4.4}$$

checking that all terms in the right hand side are in $\mathrm{Dom} D^\Lambda$ and using the induction hypothesis.

For a general $\phi \in L^2\big(S_{n+1}(\Theta), (\lambda\otimes\nu)^{n+1}\big)$ with the conditions of the theorem, first extend this function by symmetry to $([0,T]\times\Theta)^n$ Since Θ is a bounded set by hypothesis, using a mollifier family $m_\varepsilon \in \mathcal{C}_0^\infty(\mathbb{R}^{n+1})$, define

$$\phi_\varepsilon(t_1,x_1;\ldots;t_{n+1},x_{n+1}) = \int_{\mathbb{R}^{n+1}} \phi(t_1,y_1;\ldots;t_{n+1},y_{n+1}) m_\varepsilon(x_1-y_1,\ldots,x_{n+1}-y_{n+1}) \\ d\nu(y_1)\cdots d\nu(y_{n+1}).$$

Then, $\phi_\varepsilon \in \mathcal{C}_0^\infty(\mathbb{R}^{2n+2})$, $\lim_{\varepsilon\downarrow 0}\phi_\varepsilon = \phi$, uniformly and $\lim_{\varepsilon\downarrow 0}\partial_{t_j}\phi_\varepsilon = \partial_{t_j}\phi$, uniformly. Hence, by Corollary 4.2, $\lim_{\varepsilon\downarrow 0} J_{n+1}(\phi_\varepsilon) = J_{n+1}(\phi)$ in $L^2(\Omega)$, and also the right hand side of (4.3) corresponding to ϕ_ε converges in $L^2([0,T]\times\Omega)$ to the corresponding to ϕ. Thus, it suffices to prove the formula (4.3) for $\phi \in \mathcal{C}_0^\infty(\mathbb{R}^{2n+2})$. For such ϕ we can apply Theorem 1, page 65 of Yosida [29], that says that we can approximate ϕ by finite sums $\sum_{j=1}^k \phi_n^{(j)}\phi_1^{(j)}$, where $\phi_n^{(j)} \in \mathcal{C}_0^\infty(\mathbb{R}^{2n})$ and $\phi_1^{(j)} \in \mathcal{C}_0^\infty(\mathbb{R}^2)$, and also the derivatives of the sums converge uniformly to the derivatives of ϕ. Indeed, that theorem by Yosida [29] shows the result for complex valued functions, however, it can be applied to real values functions and, in that case, the approximating functions are also real. From Corollary 4.2, the symmetrization property of Remark 4.4 and the previous induction step, the proof is complete. □

Corollary 4.8 *Let Θ, Λ and ϕ be as in Theorem 4.7. With the previous notations,*

$$\phi(T_1,\Delta X_{T_1};\ldots,T_n,\Delta X_{T_n})\mathbf{1}_{\{N_T^\Theta\ge n\}} \in \mathrm{Dom}\, D^\Lambda$$

and, over $\{N_T^\Theta \ge n\}$,

$$D_t^\Lambda \phi(T_1,\Delta X_{T_1};\ldots;T_n,\Delta X_{T_n}) = \sum_{j=1}^n k(T_j,\Delta X_{T_j})\partial_j\phi(T_1,\Delta X_{T_1};\ldots,T_n,\Delta X_{T_n})\Big(\frac{T_j}{T} - \mathbf{1}_{[0,T_j]}(t)\Big).$$

Proof: We only need to observe that

$$\phi(T_1,\Delta X_{T_1};\ldots,T_n,\Delta X_{T_n})\mathbf{1}_{\{N_T^\Lambda\ge n\}} = \sum_{\ell=n}^\infty J_\ell(\psi_\ell)\mathbf{1}_{\{N_T^\Lambda=\ell\}},$$

where

$$\psi_\ell(s_1,y_1;\ldots;s_\ell,y_\ell) = \phi(s_1,y_1;\ldots;s_n,y_n), \quad \ell\ge n,$$

and apply Corollary 4.6 and Theorem 4.7. □

Remark. Note that this result yields that $D^{\{1\}}$, with $k=1$, agrees with the operator introduced in Carlen and Pardoux [8] for the Poisson process. Consequently, this paper extends the approach given in [8] to Lévy processes. Also, in general, with the previous hypothesis, the jump time T_j is in the domain of D^Λ and

$$D_t^\Lambda T_j = k(T_j,\Delta X_{T_j})\left(\frac{T_j}{T} - \mathbf{1}_{[0,T_j]}(t)\right).$$

5 Absolutely continuity for stochastic differential equations

In this section we use Proposition 3.11 and Theorem 4.7 to find conditions that guarantee that the solution at time T of some stochastic differential equations driven by a Lévy process has density with respect to the Lebesgue measure. The key point is that we can choose the convenient set Λ and weight $k(t,x)$ for each type of equation.

5.1 An equation driven by a Lévy process with continuous part

Here we consider the solution of the stochastic differential equation with an additive jump noise and a Wiener stochastic integral of the form

$$Z_t = x_0 + \int_0^t b(Z_s)ds + \int_0^t \sigma(Z_s)dW_s + \int_{(0,t]\times\mathbb{R}_0} l(y)y dN(s,y), \quad t \in [0,T],$$

where $l \in \mathcal{K}$ and the coefficients b and σ are differentiable on $\mathbb{R}$ with bounded derivatives. Then, choosing $\Lambda = \{0\}$ and $k \in \mathcal{K}$, and proceeding as in Nualart [22] (Section 2.3), we can see that

$$D_t^{\{0\}} Z_T = \sigma(Z_t) \exp\left(\int_0^t \sigma'(Z_s)dW_s + \int_0^t \left\{ b'(Z_s) - \frac{1}{2}(\sigma'(Z_s))^2 \right\} ds \right), \quad t \in [0,T].$$

Thus we can state the following theorem:

Theorem 5.1 *The random variable Z_T is absolutely continuous with respect to the Lebesgue measure on the set $\{S < T\}$, where*

$$S = \inf\left\{ t \in [0,T] : \int_0^t \mathbf{1}_{\{\sigma(Z_s)\neq 0\}} ds > 0 \right\} \wedge T.$$

Remark As in [22], we can see that last theorem also holds in the case that the coefficients are only Lipschitz functions with linear growth.

5.2 A pure discontinuous equation with a monotone drift

In this section we consider the following equation with an additive jump noise

$$Z_t = x + \int_0^t f(Z_s)ds + \int_{[0,t]\times\mathbb{R}_0} h(y)dN(s,y), \quad t \in [0,T], \tag{5.1}$$

where $x \in \mathbb{R}$, $f : \mathbb{R} \to \mathbb{R}$ is a function with a bounded derivative and $h \in L^2(\mathbb{R}_0,\nu) \cap L^1(\mathbb{R}_0,\nu)$. It is well-known that equation (5.1) has a unique square-integrable solution.

In order to calculate $D^{\mathbb{R}_0} Z_T$, we need the following result.

Proposition 5.2 *Let $\Theta_m = \{x \in \mathbb{R} : \ 1/m < |x| < m\}$, and*

$$Z_t^{(m)} = x + \int_0^t f(Z_s^{(m)})ds + \int_{[0,t]\times\Theta_m} h(y)dN(s,y), \quad t \in [0,T]. \tag{5.2}$$

Then, as $m \to \infty$, $Z_t^{(m)}$ converges to Z_t in $L^2(\Omega)$, for every $t \in [0,T]$. That convergence is also a.s. for every $t \in [0,T]$ a.e.

Proof: Let $M > 0$ be a bound for the derivative of the function f. Then

$$\mathbb{E}\left(\left|Z_t^{(m)} - Z_t\right|^2\right) \leq 2M^2T\int_0^t \mathbb{E}\left(\left|Z_s^{(m)} - Z_s\right|^2\right) ds + 2\mathbb{E}\left(\left|\int_0^t \int_{[0,t]\times(\mathbb{R}_0-\Theta_m)} h(y)dN(s,y)\right|^2\right).$$

The result is a consequence of Gronwall's lemma and of the fact that

$$\begin{aligned} &\mathbb{E}\left(\left|\int_{[0,t]\times(\mathbb{R}_0-\Theta_m)} h(y)dN(s,y)\right|^2\right) \\ &= \mathbb{E}\left(\left|\int_{[0,t]\times(\mathbb{R}_0-\Theta_m)} h(y)d\tilde{N}(s,y) + \int_0^t\int_{\mathbb{R}_0-\Theta_m} h(y)\nu(dy)ds\right|^2\right) \\ &\leq 2\int_0^T\int_{\mathbb{R}_0-\Theta_m} h(y)^2\nu(dy)ds + 2\left(\int_0^T\int_{\mathbb{R}_0-\Theta_m} |h(y)|\nu(dy)ds\right)^2 \end{aligned}$$

goes to zero as $m \to \infty$ because $\Theta_m \uparrow \mathbb{R}_0$. Finally the second claim of this result follows using a similar argument. Thus, the proof is complete. $\Box$

Now we follow a similar reasoning as Carlen and Pardoux [8]. Consider the flow $\{\Phi_t(s,x) : 0 \leq s \leq t \leq T \text{ and } x \in \mathbb{R}\}$ associated with equation (5.1). It means, $\Phi_t(s,x)$ is the unique solution to the equation

$$\Phi_t(s,x) = x + \int_s^t f(\Phi_u(s,x))du, \quad t \in [s,T].$$

Note that the partial derivatives of the flow with respect to time is

$$\partial_t\Phi_t(s,x) = f(\Phi_t(s,x)).$$

The partial derivative with respect the initial condition satisfies the equation

$$\partial_x\Phi_t(s,x) = 1 + \int_s^t f'(\Phi_u(s,x))\partial_x\Phi_u(s,x)du$$

and so

$$\partial_x\Phi_t(s,x) = \exp\left(\int_s^t f'(\Phi_u(s,x))du\right).$$

Similarly we have

$$\partial_s\Phi_t(s,x) = -f(\Phi_s(s,x)) + \int_s^t f'(\Phi_u(s,x))\partial_s\Phi_u(s,x)du.$$

Hence

$$\partial_s\Phi_t(s,x) = -f(x)\exp\left(\int_s^t f'(\Phi_u(s,x))du\right).$$

Denote by $X = \{X_t, t \geq 0\}$ the Lévy process associated to the Poisson measure N as in (2.1) (obviously with $\gamma = \sigma = 0$). The processes X and Z jump at the same times, and the jumps height of the solution process Z in a jump time τ is $h(\Delta X_\tau)$. Denote by $T_1 < T_2 < ..$ the jump times of N^{Θ_m}

(the dependence of T_j on m is suppressed to short the notations). The solution of (5.2) in $\{N_T^{\Theta_m} = n\}$ is given by

$$\begin{aligned} Z_t^{(m)} &= \Phi_t(0,x), \quad t \in [0,T_1), \\ Z_{T_1}^{(m)} &= \Phi_{T_1}(0,x) + h(\Delta X_{T_1}), \\ Z_t^{(m)} &= \Phi_t(T_1, Z_{T_1}^{(m)}), \quad t \in [T_1,T_2), \end{aligned}$$

and so on. See Figure 1.



Figure 1. A trajectory of $Z_t^{(m)}$.

By Corollary 4.8, and induction over n

$$\begin{aligned} & D_t^{\mathbb{R}_0} Z_{T_n}^{(m)} \\ = \; & f(Z_{T_n-}^{(m)}) k(T_n, \Delta X_{T_n}) \left(\frac{T_n}{T} - \mathbf{1}_{[0,T_n]}(t) \right) \\ & + \sum_{i=1}^{n-1} \exp\left(\int_{T_i}^{T_n} f'(Z_u^{(m)}) du \right) \left(f(Z_{T_i-}^{(m)}) - f(Z_{T_i}^{(m)}) \right) k(T_i, \Delta X_{T_i}) \left(\frac{T_i}{T} - \mathbf{1}_{[0,T_i]}(t) \right) \end{aligned}$$

on the set $\left\{ N_T^{\Theta_m} = n \right\}$. Thus we have that

$$\begin{aligned} & D_t^{\mathbb{R}_0} \left(Z_T^{(m)} \mathbf{1}_{\{N_T^{\Theta_m}=n\}} \right) \\ = \; & \mathbf{1}_{\{N_T^{\Theta_m}=n\}} \left[-f(Z_{T_n}^{(m)}) \exp\left(\int_{T_n}^{T} f'(Z_u^{(m)}) du \right) k(T_n, \Delta X_{T_n}) \left(\frac{T_n}{T} - \mathbf{1}_{[0,T_n]}(t) \right) \right. \\ & \left. + \exp\left(\int_{T_n}^{T} f'(Z_u^{(m)}) du \right) D_t^{\mathbb{R}_0} Z_{T_n}^{(m)} \right] \\ = \; & \sum_{i=1}^{n} \exp\left(\int_{T_i}^{T} f'(Z_u^{(m)}) du \right) \left(f(Z_{T_i-}^{(m)}) - f(Z_{T_i}^{(m)}) \right) \\ & \qquad \times k(T_i, \Delta X_{T_i}) \left(\frac{T_i}{T} - \mathbf{1}_{[0,T_i]}(t) \right) \mathbf{1}_{\{N_T^{\Theta_m}=n\}} \\ = \; & \mathbf{1}_{\{N_T^{\Theta_m}=n\}} \int_{(0,T] \times \Theta_m} \exp\left(\int_s^T f'(Z_u^{(m)}) du \right) \\ & \qquad \times \left(f(Z_{s-}^{(m)}) - f(Z_s^{\Lambda_m}) \right) k(s,y) \left(\frac{s}{T} - \mathbf{1}_{[0,s]}(t) \right) dN(s,y). \end{aligned} \tag{5.3}$$

Now we can see that the random variable Z_T is in the domain of $D^{\mathbb{R}_0,k}$.

Proposition 5.3 *Under the assumption of this section, we have that $Z_T \in \operatorname{Dom} D^{\mathbb{R}_0}$. Moreover,*

$$D_t^{\mathbb{R}_0} Z_T = \int_{[0,T]\times\mathbb{R}_0} \exp\left(\int_s^T f'(Z_u)du\right)(f(Z_{s-}) - f(Z_s))\,k(s,y)\left(\frac{s}{T} - \mathbf{1}_{[0,s]}(t)\right)dN(s,y). \quad (5.4)$$

Proof: The result follows from Proposition 5.2, letting $m \to \infty$ and from the dominated convergence theorem. That theorem can be applied due to the integrand in the right-hand side of (5.3) is bounded by $M|h(y)k(s,y)|$, for some $M > 0$, since f' is bounded, and the jumps of the solution are related with h. $\square$

5.2.1 Case with finite Lévy measure

In this section we analyze the existence of a density for Z_T in the case that the Lévy measure is finite and the drift is a monotone function.

Theorem 5.4 *Assume that $\nu(\mathbb{R}_0) < \infty$. Consider equation (5.1), where f is a momotone function with bounded derivative and $h \in L^2(\mathbb{R}_0,\nu))$ with $h(y) \neq 0$ for $y \in \mathbb{R}_0$. Then Z_T is absolutely continuous on the set $\left\{N_T^{\mathbb{R}_0} \geq 1\right\}$.*

Proof: We first assume that f is an increasing function. We choose for k the function $(-h \wedge 1) \vee (-1)$ that depends only on y. Then, $(f(Z_{s-}) - f(Z_{s-} + h(\Delta X_s)))k(s,\Delta X_s)$ is strictly positive in the jump points. Thus, to finish the proof, we only need to apply Proposition 3.11 and observe that

$$\int_0^T \left(\frac{s}{T} - \mathbf{1}_{[0,s]}(t)\right)^2 dt = s(1-\frac{s}{T}), \quad s \in [0,T],$$

and

$$\int_0^T \left(\frac{s}{T} - \mathbf{1}_{[0,s]}(t)\right)\left(\frac{r}{T} - \mathbf{1}_{[0,r]}(t)\right)dt = s(1-\frac{r}{T}), \quad \text{for } 0 \leq s < r \leq T.$$

Hence the two integrals are positive for $s < t$. Indeed, we have that

$$\begin{aligned}
&\int_0^T \left(D_t^{\mathbb{R}_0} Z_T\right)^2 dt \\
&= \sum_{i=1}^{N_T^{\mathbb{R}_0}} \exp\left(2\int_{T_i^{\mathbb{R}_0}}^T f'(Z_u)du\right)\left(f(Z_{T_i^{\mathbb{R}_0}}-) - f(Z_{T_i^{\mathbb{R}_0}})\right)^2 (k(\Delta X_{T_i^{\mathbb{R}_0}}))^2 T_i^{\mathbb{R}_0}\left(1 - \frac{T_i^{\mathbb{R}_0}}{T}\right) \\
&\quad +2\sum_{1\leq i<j\leq N_T^{\mathbb{R}_0}} \exp\left(\int_{T_i^{\mathbb{R}_0}}^T f'(Z_u)du\right)\left(f(Z_{T_i^{\mathbb{R}_0}}-) - f(Z_{T_i^{\mathbb{R}_0}})\right)k(\Delta X_{T_i^{\mathbb{R}_0}}) \\
&\qquad \times \exp\left(\int_{T_j^{\mathbb{R}_0}}^T f'(Z_u)du\right)\left(f(Z_{T_j^{\mathbb{R}_0}}-) - f(Z_{T_j^{\mathbb{R}_0}})\right)k(\Delta X_{T_j^{\mathbb{R}_0}}) T_i^{\mathbb{R}_0}\left(1 - \frac{T_j^{\mathbb{R}_0}}{T}\right),
\end{aligned}$$

which is bigger that zero on the set $\left\{N_T^{\mathbb{R}_0} \geq 1\right\}$. Therefore Z_T absolutely continuous in this set.

Finally we can proceed similarly in the case that f is decreasing using the function $k = (h \wedge 1) \vee (-1)$. $\square$

5.2.2 Case with infinite Lévy measure

Now we deal with the case that f is only monotone on an neighborhood of the initial condition x. This problem has been analyzed in Nourdin and Simon [21] using a stratification method.

Theorem 5.5 *Consider equation (5.1), where f is a function with a bounded derivative and $h \in L^2(\mathbb{R}_0, \nu) \cap L^1(\mathbb{R}_0, \nu)$. Assume that $\nu(\mathbb{R}_0) = \infty$, $h(y) \neq 0$ for $y \in \mathbb{R}_0$, and that f is monotone on a neighborhood of the point x. Then, the random variable Z_T is absolutely continuous.*

Proof: Here it is convenient to write the function k that we use in the derivative, $D^{\Lambda,k}$. Proceeding as in Theorem 5.4, we assume that there is $\varepsilon > 0$ such that f is increasing on $(x-\varepsilon, x+\varepsilon)$; the proof for f decreasing is similar.

Note that (5.1), together with the fact that f has a bounded derivative, implies that there exists $M > 0$ such that

$$|Z_t - x| \leq M \int_0^t |Z_s - x| ds + t|f(x)| + \int_{[0,t]\times\mathbb{R}_0} |h(y)| dN(s,y), \quad t \in [0,T].$$

Consequently, Gronwall's lemma leads to write

$$|Z_t - x| \leq \left(t|f(x)| + \int_{[0,t]\times\mathbb{R}_0} |h(y)| dN(s,y) \right) e^{MT}, \quad t \in [0,T].$$

Therefore, there is $t_0 \in (0,T)$ such that

$$|Z_t - x| \leq \frac{\varepsilon}{2} + e^{MT} \int_{[0,t]\times\mathbb{R}_0} |h(y)| dN(s,y), \quad t \in [0,t_0]. \tag{5.5}$$

Now, for $t \in [0,t_0]$, let

$$A_t = \left\{ e^{MT} \int_0^t \int_{\mathbb{R}_0} |h(y)| dN(s,y) > \frac{\varepsilon}{2} \right\}$$

and $k_t : [0,T] \to \mathbb{R}$ a function in $C^1((0,T))$ such that $k_t(s) > 0$ for $s \in [0,t)$, and $k_t(s) = 0$ for $s \geq t$. Define $k^{(t)}(s,y) = -k_t(s)((h(y) \wedge 1) \vee (-1))$. As the Lévy measure is ∞, there are, with probability 1, jumps in $(0,t)$, and from (5.5) we deduce that the jump sizes satisfy $|Z_s - x| \leq \varepsilon$ for $s \leq t \leq t_0$ on the set A_t^c.

Indeed, in this case we have that, on A_t^c, $(f(Z_{s-}) - f(Z_s))k^{(t)}(s,y) \geq 0, \quad$ for $0 \leq s \leq T$, with $(f(Z_{s-}) - f(Z_s))k^{(t)}(s,y) > 0$ if Z has a jump at time $s \in (0,t)$. So, (5.4) gives

$$\int_0^T \left(D_u^{\mathbb{R}_0, k^{(t)}} Z_T \right)^2 du > 0 \quad \text{in } A_t^c.$$

As a consequence, by Proposition 3.11, the condition $\lambda(B) = 0$ implies that

$$P(\{Z_T \in B\} \cap A_t^c) = 0.$$

Finally, we observe that $A_t^c \subset A_{t'}^c$ for $t' < t$, and using the Markov inequality, we have

$$\begin{aligned} P(A_t) &\leq \frac{4e^{2MT}}{\varepsilon^2} E\left(\left(\int_{[0,t]\times\mathbb{R}_0} |h(y)| dN(s,y) \right)^2 \right) \\ &\leq \frac{8e^{2MT}}{\varepsilon^2} t \left(\int_{\mathbb{R}_0} |h(y)|^2 \nu(dy) + T \left(\int_{\mathbb{R}_0} |h(y)| \nu(dy) \right)^2 \right). \end{aligned}$$

Hence we see that $\lim_{t\downarrow 0} P(A_t) = 0$.

So $P(\cup_{t<t_0,\, t\in\mathbb{Q}} A_t^c) = 1$, and we can conclude, as $P(\{Z_T \in B\}) = \lim_{t\downarrow 0} P(\{Z_T \in B\} \cap A_t^c)$, that Z_T is absolutely continuous. $\square$

5.3 A pure discontinuous equation with non zero Wronskian

In this section we assume that the Lévy measure ν is finite and consider the stochastic differential equation

$$Z_t = x + \int_0^t f(Z_s)ds + \int_0^t \int_{\mathbb{R}_0} h(y)g(Z_{s-})dN(s,y), \quad t \in [0,T], \tag{5.6}$$

where $x \in \mathbb{R}$, $h \neq 0$ is a bounded function in $L^2(\mathbb{R}_0, \nu)$, and $f : \mathbb{R} \to \mathbb{R}$ and $g : \mathbb{R} \to \mathbb{R}$ are two bounded functions with two bounded derivatives and one bounded derivative, respectively. The existence of a density for Z_T was analyzed by Carlen and Pardoux [8] in the case that the involved Lévy process is a Poisson process. In the remaining of this section we also assume that

$$|h(y)W(g,f)(x)| > \frac{1}{2}||f''||_\infty ||h||_\infty^2 ||g||_\infty^2, \quad x \in \mathbb{R}\ y \in \mathbb{R}_0, \tag{5.7}$$

where $W(g,f) = g'f - f'g$ is the Wronskian of g and f.

As in Section 5.2, we consider the flow $\{\Phi_t(s,x) : 0 \leq s \leq t \leq T \text{ and } x \in \mathbb{R}\}$ associated with equation (5.6) and the family of jump times $\{T_j : j \in \mathbb{N}\}$ of the Lévy process X. Remember that now we are dealing with a finite Lévy measure in this part of the paper.

Note that the fact that $Z_{T_n} = \Phi_{T_n}(T_{n-1}, Z_{T_{n-1}}) + h(\Delta X_{T_n})g(Z_{T_n -})$ jointly with $T_0 = 0$ and Corollary 4.8 allow to utilize induction on n to see that, for any $k \in \mathcal{K}$,

$$\begin{aligned} D_t^{\mathbb{R}_0} Z_{T_n} &= f(Z_{T_n -})k(T_n, \Delta X_{T_n})\left(\frac{T_n}{T} - \mathbf{1}_{[0,T_n]}(t)\right) \\ &\quad + f(Z_{T_n -})h(\Delta X_{T_n})g'(Z_{T_n -})k(T_n, \Delta X_{T_n})\left(\frac{T_n}{T} - \mathbf{1}_{[0,T_n]}(t)\right) \\ &\quad + \sum_{i<n} \exp\left(\int_{T_i}^{T_n} f'(Z_u)du\right) r_t(T_i, \Delta X_{T_i}) \\ &\quad + \sum_{m=2}^{n} \int_{S_m(T_m)} \exp\left(\int_{s_1}^{T_n} f'(Z_u)du\right) h(y_m)g'(Z_{s_m -}) \cdots h(y_2)g'(Z_{s_2 -}) r_t(s_1, y_1) \\ &\qquad \times dN(s_m, y_m) \cdots dN(s_1, y_1), \end{aligned} \tag{5.8}$$

where

$$r_t(s,y) = k(s,y)\left(\frac{s}{T} - \mathbf{1}_{[0,s]}(t)\right)\left(f(Z_{s-}) - f(Z_s) + f(Z_{s-})h(y)g'(Z_{s-})\right)$$

and

$$S_m(r) = \{(s_1, y_1; \ldots; s_m, y_m) \in ([0,T] \times \mathbb{R}_0)^m : 0 \leq s_1 < \cdots < s_m \leq r\}.$$

Now we are ready to see that Z_T is in the domain of $D^{\mathbb{R}_0}$.

Proposition 5.6 *Let Z be the solution of equation (5.6). Then $Z_T \in \operatorname{Dom} D^{\mathbb{R}_0}$.*

Proof: By Corollary 4.8 and (5.8), we have $Z_T\mathbf{1}_{[N_T^{\mathbb{R}_0}=n]}$ is in the domain of $D^{\mathbb{R}_0}$ and

$$\begin{aligned}
D_t^{\mathbb{R}_0}(Z_T\mathbf{1}_{[N_T^{\mathbb{R}_0}=n]}) &= D_t^{\mathbb{R}_0}\left(\Phi_T(T_n, Z_{T_n})\mathbf{1}_{\{N_T^{\mathbb{R}_0}=n\}}\right)\\
&= \mathbf{1}_{[N_T^{\mathbb{R}_0}=n]}\left\{-f(Z_{T_n})\exp\left(\int_{T_n}^T f'(Z_u)du\right)k(T_n,\Delta X_{T_n})\left(\frac{T_n}{T}-\mathbf{1}_{[0,T_n]}(t)\right)\right.\\
&\quad \left. + \exp\left(\int_{T_n}^T f'(Z_u)du\right)D^{\mathbb{R}_0}(Z_{T_n})\right\}\\
&= \mathbf{1}_{[N_T^{\mathbb{R}_0}=n]}\left\{\sum_{i\le n}\exp\left(\int_{T_i}^T f'(Z_u)du\right)r_t(T_i,\Delta X_{T_i})\right.\\
&\quad +\sum_{m=2}^n\int_{S_m(T_n)}\exp\left(\int_{s_1}^T f'(Z_u)du\right)k(y_m)g'(Z_{s_m-})\cdots h(y_2)g'(Z_{s_2-})r_t(s_1,y_1)\\
&\qquad \left.\times\, dN(s_1,y_1)\cdots dN(s_m,y_m)\right\}.
\end{aligned}$$

Hence, the fact that there is a constant $C>0$ such that $|r_t(s,y)|<Ck(s,y)$ and

$$\begin{aligned}
&\left|\int_{S_m(T_n)}\exp\left(\int_{s_1}^T f'(Z_u)du\right)\right.\\
&\qquad\qquad \left.\times\, h(y_m)g'(Z_{s_m-})\cdots h(y_2)g'(Z_{s_2-})r_t(s_1,y_1)dN(s_1,y_1)\cdots dN(s_m,y_m)\right|\\
&\le \mathbf{1}_{\{m\le n\}}\frac{C}{m!}\left(\int_0^T\int_{\mathbb{R}_0}|k(s,y)|\,dN(s,y)\right)\left(\int_0^T\int_{\mathbb{R}_0}|h(y)|\,dN(s,y)\right)^{m-1}
\end{aligned}$$

implies that Z_T is in the domain of $D^{\mathbb{R}_0}$ with

$$\begin{aligned}
D_t^{\mathbb{R}_0}Z_T &= \int_{[0,T]\times\mathbb{R}_0}\exp\left(\int_s^T f'(Z_u)du\right)r_t(s,y)dN(s,y)\\
&\quad +\sum_{m=2}^\infty\int_{S_m(T)}\exp\left(\int_{s_1}^T f'(Z_u)du\right)h(y_m)g'(Z_{s_m-})\cdots h(y_2)g'(Z_{s_2-})r_t(s_1,y_1)\\
&\quad \times dN(s_1,y_1)\cdots dN(s_m,y_m). \qquad (5.9)
\end{aligned}$$

Thus the proof is complete. $\square$

Finally we can see that Z_T has a density.

Theorem 5.7 *Let Z be the solution of equation (5.6). Then Z_T has a density on the set $[N_T^{\mathbb{R}_0}>0]$.*

Proof: For a rational number $p\in(0,T)$ and a positive integer $n\ge 1$, we introduce the set

$$A_{p,n}=\{N_T=n\}\cap\{T_{n-1}\le p<T_n\},$$

with the convention $T_0=0$. We choose a function $h_p:[0,T]\to\mathbb{R}_+$ of class $C^1((0,T))$ such that $h_p(s)=0$ for $s\le p$, and $h_p(s)>0$ for $s>p$. Then, (5.9) implies that in the set $A_{p,n}$ we only have to consider in the derivative the jump at T_n,

$$\begin{aligned}
D_t^{\mathbb{R}_0,h_p}Z_T &= \exp\left(\int_{T_n}^T f'(Z_u)du\right)h_p(T_n)\left(\frac{T_n}{T}-\mathbf{1}_{[0,T_n]}(t)\right)\\
&\qquad\times\left(f(Z_{T_n-})-f(Z_{T_n})+f(Z_{T_n-})h(\Delta X_{T_n})g'(Z_{T_n-})\right),
\end{aligned}$$

which is different than zero due to condition (5.7). Hence Proposition 3.11 implies that $P(\{Z_T \in B\} \cap A_{p,n}) = 0$ for any Borel set $B \in \mathcal{B}(\mathbb{R})$ such that $\lambda(B) = 0$. Thus, for this Borel set we have

$$P(\{Z_T \in B\} \cap \{N_T > 0\}) = P(\{Z_T \in B\} \cap (\cup_{n\in\mathbb{N},p\in\mathbb{Q}} A_{p,n})) = 0,$$

and the proof is finished. $\square$

Acknowledgment. The authors thank Cinvestav-IPN, Universitat Autònoma de Barcelona and Universitat de Barcelona for their hospitality and economical support during the realization of this paper.

References

[1] R.A. Adams (1975) *Sobolev Spaces.* Pure and Applied Mathematics, Vol. **65**. Academic Press.

[2] D. Applebaum (2009) Universal Malliavin calculus in Fock and Lévy-Ito spaces. *Communications in Stoch. Anal.* **3**(1), pp. 119-141.

[3] V. Bally (2007) Integration by parts formula for locally smooth laws and applications to equations with jumps I and II. *Preprints of Institut Mittag-Leffler* No. 14-15.

[4] F.E. Benth and A. Lokka (2004) Anticipative calculus for Lévy processes and stochastic differential equations. *Stoch. and Stoch. Reports* **76**(3), pp. 191-211.

[5] K. Bichteler, J.B. Gravereaux and J. Jacod (1987) *Malliavin Calculus for Processes with Jumps*. Stochastics Monographs **2**, Gordon and Breach Science Publishers.

[6] J.M. Bismut (1983) Calcul des variations stochastique et processus de sauts. *Z. Wahrsch. verw. Gebiete* **63**(2), pp. 147-235.

[7] N. Bouleau and L. Denis (2009) Energy image density property and the lent particle method for Poisson measures. *J. Funct. Anal.* **257**(4), pp. 1144-1174.

[8] E.A. Carlen and E. Pardoux (1990) Differential calculus and integration by parts on a Poisson space. In *Stochastics, Algebra and Analysis in Classical and Quantum Dynamics, S. Albeverio et al. (eds)*, pp. 63-73. Kluwer.

[9] Y.A. Davydov, M.A. Lifshits and N.V. Smorodina (1998) Local Properties of distributions of stochastic functionals. *Translations of Mathematical Monographs* **173**, AMS, pp. 509-548.

[10] L. Decreusefond and N. Savy (2006) Anticipative calculus with respect to filtered Poisson processes. *Ann. Inst. H. Poincaré Probab. Statist* **42**(3), pp. 343-372.

[11] L. Denis (2000) A criterion of density for solutions of Poisson-driven SDEs. *Probab. Theory Related Fields* **118**(3), pp. 406-426.

[12] G. Di Nunno (2002) Stochastic integral representations, stochastic derivatives and minimal variance hedging. *Stoch. Stoch. Rep.* **73**(1-2), pp. 181-198.

[13] G. Di Nunno, T. Meyer-Brandis, B. Oksendal and F. Proske (2005) Malliavin calculus and anticipative Itô formulae for Lévy processes. *Infinite Dimensional Anaysis, Quantum Probability and Related Topics* **8**(2), pp. 235-258.

[14] C. Geiss and E. Laukkarinen (2011) Denseness of certain smooth Lévy functionals in $\mathbb{D}_{1,2}$. *Probab. Math. Statist.* **31**(1), pp. 1-15

[15] Y. Ishikawa and H. Kunita (2006) Malliavin calculus on the Wiener-Poisson space and its application to canonical SDE with jumps. *Stochastic Process. Appl.* **116**(12), pp. 1743-1769.

[16] K. Itô (1956) Spectral type of shift transformation of differential processes with stationary increments. *Trans. Amer. Math. Soc.* **81**, pp. 253-263.

[17] A.M. Kulik (2011) Absolute continuity and convergence in variation for distributions of a functionals of Poisson point measure. *J. Theoret. Probab.* **24**(1), pp. 1-38.

[18] R. Léandre (1985) Régularité de processus de sauts dégénérés. *Ann. Ins. H. Poincaré. Probab. Statist.* **21**(2), pp. 125-146.

[19] Y.J. Lee and H.H. Shih (2004) The product formula of multiple Lévy-Itô integrals. *Bull. Inst. Math. Acad. Sinica* **32**(2), pp. 71-95.

[20] J.A. León and C. Tudor (1998) A chaos approach to anticipanting calculus for the Poisson process. *Stochastics Stochastics Rep.* **62**(3-4), pp. 217-250.

[21] I. Nourdin and T. Simon (2006) On the absolute continuity of Lévy processes with drift. *The Annals of Probability* **34**(3), pp. 1035-1051.

[22] D. Nualart (2006) *Malliavin Calculus and Related Topics. Second Edition.* Springer.

[23] D. Nualart and J. Vives (1990) Anticipative calculus for the Poisson process based on the Fock space. *Séminaire de Probabilités* **XXIV**, LNM 1426, pp. 154-165.

[24] J. Picard (1996) On the existence of smooth densities for jump processes. *Probab. Theory Related Fields* **105**(4), pp. 481-511.

[25] J. Picard (1996) Formules de dualité sur l'espace de Poisson. *Ann. Inst. Henri Poincaré* **32**(4), pp. 509-548.

[26] J.Ll. Solé, F. Utzet and J. Vives (2007) Canonical Lévy process and Malliavin calculus. *Stochastic Process. Appl.* **117**(2), 165-187.

[27] K.I. Sato (1999) *Lévy Processes and Infinitely Divisible Distributions.* Cambridge Studies in Advanced Mathematics **68**. Cambridge.

[28] C. Tudor (1997) A product formula for multiple Poisson-Itô integrals. *Revue Roum. Math. Pures et Appl.* **XLII**(3-4), 339-345.

[29] K. Yosida (1980) *Functional Analysis.* Sixth Edition. Springer.